

## First record of *Metopograpsus cannicci* Innocenti, Schubart and Fratini, 2020 (Crustacea: Decapoda: Grapsidae) from India

Mithila Bhat<sup>1</sup>, Krupal J Patel<sup>2</sup> and Jigneshkumar N Trivedi<sup>3\*</sup>

<sup>1</sup>School of Earth, Ocean and Atmospheric Sciences, Goa University, Taleigao-403206, Goa, India

<sup>2</sup>Marine Biodiversity and Ecology Laboratory, Department of Zoology, The Maharaja Sayajirao University of Baroda, Vadodara-390002, Gujarat, India

<sup>3</sup>Department of Life Sciences, Hemchandracharya North Gujarat University, Patan-384265, Gujarat, India

\*Corresponding author ✉: [jntrivedi26@yahoo.co.in](mailto:jntrivedi26@yahoo.co.in)

### Abstract

Received: 25 August 2021

Accepted: 19 December 2021

Published online: 31 December 2021

*Metopograpsus cannicci* Innocenti, Schubart and Fratini, 2020 is distributed in the Red Sea, Iran, along the East African coast, Seychelles, Madagascar, and Mauritius. The present paper reports this species for the first time from India indicating its eastward extension.

**Key words:** Brachyura, mangroves, new record, rocky shores, West coast of India

### Introduction

The genus *Metopograpsus* H. Milne Edwards, 1835 comprises seven species: *M. messor* (Forskål, 1775); *M. thukuhar* (Owen, 1839); *M. oceanicus* (Hombron and Jacquinot, 1846 [in Hombron and Jacquinot, 1842–1854]); *M. latifrons* (White, 1847); *M. quadridentatus* Stimpson, 1858; *M. frontalis* Miers, 1880 and *M. cannicci* Innocenti, Schubart and Fratini, 2020. The taxonomy of *Metopograpsus* is difficult owing to the subtle differences in the morphological features between species (Fratini et al., 2018; Innocenti et al., 2020). Recently Innocenti et al. (2020) examined specimens identified as *M. thukuhar* and on the basis of their morphological and genetic investigations described a new species, *M. cannicci*, referring to it as a pseudocryptic taxon.

Fratini et al. (2018), through molecular phylogenetic investigation found that the hitherto widely distributed *M. thukuhar* comprised two species with different distributions. *M. thukuhar* is confined to Southeast Asia and Pacific islands, while populations from the East African coast, Red Sea and Seychelles were described as the distinct species, *M. cannicci* by Innocenti et al. (2020). In the present study, *M. cannicci* is reported for the first time from India.

### Material and Methods

The specimens were collected by hand-picking from the mangrove region of Goa and Gujarat states of India (Fig. 1). The specimens were washed to remove debris and photographed. They were preserved in 90% ethanol and deposited in the Zoological Reference Collection (LFSC.ZRC), Department of Life Sciences, Hemchandracharya North Gujarat University, Patan, Gujarat, India. The size of the specimens was recorded using digital vernier callipers (INSIZE, Model no. 1108200).

### Abbreviations

CW, carapace width; CL, carapace length; G1, male first gonopod; coll., collector. Morphological terminology used in this article follows Innocenti et al. (2020).

### Results and Discussion

#### Taxonomy

**Order Decapoda Latreille, 1802**

**Superfamily Grapsoidea MacLeay, 1838**

**Family Grapsidae MacLeay, 1838**

**Genus *Metopograpsus* H. Milne Edwards, 1853**

***Metopograpsus cannicci* Innocenti, Schubart and Fratini, 2020 (Fig. 2)**

*Metopograpsus thukuhar* – Crosnier, 1965: 25, Fig. 20–22, 27; Hartnoll, 1975: 317; Vannini and Valmori, 1981: 73, Fig. 8B, 9B; Naderloo, 2011: Figs. 4a–g, 5f, Figs. 5f.; Trivedi et al., 2018: 42 (in list).

*Metopograpsus cannicci* – Innocenti et al., 2020: 621, Figs. 1A, 2A, 3A, 4A, 5A, 6.

#### Material examined

7 males (CW: 13.74–25.35 mm, CL: 10.20–19.82 mm); 8 females (CW: 13.40–28.88 mm, CL: 10.00–23.14 mm) LFSC.ZRC- 36, mangroves, Chapora estuary (15°37'22.2"N, 73°44'52.3"E), Goa, India, 10 November 2016. 3 males (CW: 12.00–20.68 mm, CL: 9.30–16.20 mm); 4 females (CW: 12.40–17.49 mm, CL: 9.12–13.04 mm), LFSC.ZRC- 36, mangroves, Sal estuary (15°10'05.7"N, 73°56'46.9"E), Goa, India, 8 October 2016. 3 males (CW: 18.47–27.06 mm, CL: 14.76–22.07 mm); 3 females (CW: 19.68–27.72 mm, CL: 15.02–22.09 mm) LFSC.ZRC- 36, Rocky shore, Cakra beach

(15°27'03.9"N, 73°50'15.9"E), Goa, India, 12 February 2020. coll. M. Bhat. 9 males (CW: 10.15–29.51 mm, CL: 7.36–23.18 mm); 14 females (CW: 12.16–25.15 mm, CL: 8.37–18.24 mm) LFSC.ZRC- 36, Mohadi (23°19'56.1"N, 68°37'20.6"E), Gujarat, India, 22 May 2011. coll. J. Trivedi.


#### Species description (modified from Innocenti et al., 2020)

Carapace flat, quadrangular, broader than long, and smooth. Lateral margins entire. Regions weakly defined, urogastric region with groove distinct, branchial region having distinct oblique ridges, cardiac and intestinal regions smooth, with no ridge or tubercles (Fig. 2a). Front broad, deflexed with rugose surface and free margin crenulated, with some concavity, 4 depressed post-frontal lobes along the line of frontal deflexion. Suborbital tooth triangular, suborbital border denticulate (Fig 2d). The exposed surface of the base of the antenna densely pubescent.



**Figure 1:** Mangrove habitat inhabited by *Metopograpsus cannicci* Innocenti, Schubart and Fratini, 2020.





**Figure 2:** *Metopograpsus cannicci* Innocenti, Schubart, and Fratini 2020; male- CL 20.61 mm; CW: 25.35 mm; female- CL 19.29 mm; CW: 24.73 mm: a. dorsal habitus; b. ventral habitus; c. left chela, outer view; d. detail of suborbital region; e. male abdomen; f. first left pleopod (G1), dorsal view; g. G1 apical chitinous process; h. female gonopore.

Chelipeds subequal, with no certain handedness, fingers stout with spatulated tip, slight gape visible when closed. The cutting edge of both fingers with a series of inconspicuous teeth (Fig 2c). Ambulatory legs compressed, merus broad and longest, third and fourth pereiopod longest, first shortest.

Male abdomen with 6 distinct somites, first as broad as thoracic sternum, somite 6 almost rectangular, telson

triangular (Fig. 2b, e). Female pleon fringed with long setae, broad, evenly rounded, telson half-moon shaped.

G1 slender, with twisted shaft along the longitudinal axis, (Fig 2f) apical corneous process elongate, bearing long setae at the base (Fig, 2g). In female, vulva in depression at sternite 5/6 border, operculum on inner part with an evident trapezoidal sternal vulvar cover in adult females (Fig. 2h).

## Coloration

Small crabs (< 23 mm CW) were observed mostly in different shades of mottled green with brown, black, and cream patterns on the carapace. Larger crabs (> 23 mm CW) exhibited a mostly dark brown carapace. Ventral surface white. Fingertips of chela were bright orange. Pereiopods almost matching the color of the carapace with a pattern.

## Distribution

*Metopograpsus cannicci* has been previously recorded from Iran, Somalia, Kenya, Tanzania, Mozambique, Seychelles, Madagascar, Mauritius (Innocenti et al., 2020), and in Gujarat and Goa, India in the present study.

## Remarks

The Indian specimens generally agree with the description and figures of *M. cannicci* (Innocenti et al., 2020) but differ slightly in having the suborbital border smaller and with more closely spaced denticles than in the type specimens (Fig 2d).

*Metopograpsus cannicci* closely resembles *M. thukuhar* but can be differentiated from the latter species on the basis of the following characters: the G1 less swollen, a more acute hump at its distal end, and a longer corneous tip in *M. cannicci* (Fig. 2f) (versus G1 distally more swollen, with a rounded hump and a shorter corneous tip in *M. thukuhar*, Innocenti et al., 2020, Fig. 4B) and the adult females having a trapezoidal sternal vulvar cover in *M. cannicci* (Fig. 2h) (versus, adult females having a wider and higher cover in *M. thukuhar*, Innocenti et al., 2020, Fig. 5B).

*Metopograpsus thukuhar* has been recorded from different coastal states of India including Gujarat (Trivedi et al., 2018), Tamil Nadu (Silas and Sankarankutty, 1965; Thomas, 1969; Venkataraman et al., 2004; Dev Roy and Nandi, 2007), Odisha (Dev Roy and Rath, 2017) and Andaman and Nicobar Islands (Kumarlingam et al., 2017). We have re-examined the specimens collected and identified as *M. thukuhar* by Trivedi et al. (2018) from Gujarat and can confirm that these belong to *M. cannicci*. All other records of *M. thukuhar* from the East coast of India appear only in species checklists without mention of voucher specimens. We believe that all the records of *M. thukuhar* from India may belong to *M. cannicci*.

In the present study, *M. cannicci* was found in mangroves as well as on open rocky shore habitat. Smaller crabs were mostly found on the ground, whereas larger crabs were found mostly between crevices, holes in the tree trunks of mangroves and between rock/boulder crevices on the rocky shore. The mangrove environment was mostly made up of *Avicennia officinalis*, *Avicennia marina* and *Rhizophora mucronata*. In the present

study, *M. cannicci* was found in sympatry with *M. latifrons* (White, 1847), *Austruca annulipes* (H. Milne Edwards, 1837) and *Parasesarma bengalense* (Davie, 2003).

## Acknowledgement

The authors are thankful to two anonymous reviewers for their valuable comments on the manuscript.

## Conflict of interest

All authors declare that there are no conflicting issues related to this research article.

## References

- Crosnier, A. (1965). Crustacés Dècapodes, Grapsidae et Ocypodidae. *Faune de Madagascar*, 18: 1–14.
- Dev Roy, M. K. and Nandi, N. C. (2007). Brachyuran diversity in coastal ecosystems of Tamil Nadu. *Journal of Environment and Sociobiology*, 4 (2): 169–192.
- Dev Roy, M. K. and Rath, S. (2017). An inventory of crustacean fauna from Odisha Coast, India. *Journal of Environment and Sociobiology*, 14 (1): 49–112.
- Forskål, P. (1775). *Descriptiones Animalium, Avium, Amphibiorum, Piscium, Insectorum, Vermium; quae in Itinere Orientali Observavit Petrus Forskål. Post Mortem Auctoris editit Carsten Niebuhr*. Adjuncta est materia Medica Kahirina. Mölleri, Hafniae, 19 + xxxiv + 164 pp.
- Fratini, S., Canticci, S., and Schubart, C. D. (2018). Molecular phylogeny of the crab genus *Metopograpsus* H. Milne Edwards, 1853 (Decapoda: Brachyura: Grapsidae) reveals high intraspecific genetic variation and distinct evolutionarily significant units. *Invertebrate Systematics*, 32 (1): 215–223.  
<https://doi.org/10.1071/IS17034>
- Hartnoll, R. G. (1975). The Grapsidae and Ocypodidae (Decapoda: Brachyura) of Tanzania. *Journal of Zoology*, 177: 305–328.  
<https://doi.org/10.1111/j.1469-7998.1975.tb02235.x>
- Hombron, J. B. and Jacquinot, H. (1842–1854). *Crustacés. Atlas d'Histoire Naturelle. Zoologie. Voyage au Pôle Sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée pendant les années 1837–1838–1839–1840, Crustacés*. Gide et J. Baudry, Paris, pls. 1–9.
- Innocenti, G., Schubart, C. D. and Fratini, S. (2020). Description of *Metopograpsus cannicci*, new species, a pseudocryptic crab species from East Africa and the Western Indian Ocean (Decapoda: Brachyura: Grapsidae). *Raffles Bulletin of Zoology*, 68: 619–628.  
<https://doi.org/10.26107/RBZ-2020-0077>

- Kumaralingam, S., Raghunathan, C., Ajithkumar, T. T. and Chandra, K. (2017). Reef associated common crustaceans of Andaman and Nicobar islands, Lakshadweep and Gulf of Mannar. *Records of Zoological Survey of India, Occasional Paper*, 380: 1–249.
- Latreille, P. A. (1802). *Histoire naturelle, générale et particulière des Crustacés et des Insectes. Ouvrage faisant suite à l'histoire naturelle générale et particulière, composée par Leclerc de Buffon, et rédigée par C.S. Sonnini, membre de plusieurs sociétés savantes. Familles naturelles des genres*. Vol. 3. Paris, F. DuFart, 467 pp.
- MacLeay, W. S. (1838). On the brachyurous decapod crustacea brought from the Cape by Dr Smith, In: MacLeay, W. S. (Ed.), *Illustrations of the Zoology of South Africa; being a portion of the objects of natural history chiefly collected during an expedition into the interior of South Africa, under the direction of Dr Andrew Smith, in the Years 1834, 1835, and 1836; Fitted out by "The Cape of Good Hope Association for Exploring Central Africa"*. Smith, Elder & Co., London. pp. 53–71.
- Miers, E. J. (1880). On a Collection of Crustacea from the Malaysian Region.—Part II. Telpusidea, Catometopa, and Oxystomata. *Annals and Magazine of Natural History, series 5*. 5 (28): 304–317; Pl. XIV.
- Milne Edwards, H. (1853). Mémoire sur la famille des Ocypodides. Suite (1). Deuxième Tribu Principale. *Annales des Sciences Naturelles*, 3e série, 20: 163–228
- Naderloo, R. and Türkay, M. (2012). Decapod crustaceans of the littoral and shallow sublittoral Iranian coast of the Persian Gulf: Faunistics, biodiversity and zoogeography. *Zootaxa*, 3374 (1): 1–67. <https://doi.org/10.11646/zootaxa.3374.1.1>
- Owen, R. (1839). Crustacea, In: *The zoology of Captain Beechey's Voyage to the Pacific Ocean and Bering's Strait, performed in H.M.S. Blossom in the years 1825–1828*. London, 144 pp.
- Silas, E. G. and Sankarankutty, C. (1965). Field investigations on the shore crabs of the Gulf of Mannar and Palk bay, with special reference to the ecology and behaviour of the pellet crab *Scopimera proxima* Kemp. *Proceedings of Symposium on Crustacea*, 3: 1008–1025.
- Stimpson, W. (1858). Prodromus descriptionis animalium evertibratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars V. Crustacea Ocypodoidea. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 10: 93–110.
- Thomas, M. M. (1969). Catalogue of crabs and hermit-crabs in the reference collections of the central marine fisheries research institute. *Bulletin of the Central Marine Fisheries Research Institute*, 7: 41–50.
- Trivedi, J. N., Trivedi, D. J., Vachhrajani, K. D. and Ng, P. K. L. (2018). An annotated checklist of the marine brachyuran crabs (Crustacea: Decapoda: Brachyura) of India. *Zootaxa*, 4502 (1): 1–83. <https://doi.org/10.11646/zootaxa.4502.1.1>
- Vannini, M. and Valmori, P. (1981). Researches on the coast of somalia. The shore and the dune of sar uanle. 30. Grapsidae (Decapoda Brachyura). *Monitore Zoologico Italiano, Supplemento*, 14 (1): 57–101. <https://doi.org/10.1080/03749444.1981.10736614>
- Vekataraman, K., Jeyabaskaran, R., Raghuram, K. P. and Alfred, J. R. B. (2004). Bibliography and checklist of corals and coral reef-associated organisms of India. *Records of Zoological Survey of India, Occasional Paper*, 226: 1–468.
- White, A. (1847). No. VIII. Descriptions of a new genus and five new species of Crustacea. Appendix, In: Jukes, J. B. (Ed.), *Narrative of the surveying voyage of H.M.S. Fly, commanded by Captain F.P. Blackwood, R.N. in Torres Strait, New Guinea, and other islands of the Eastern Archipelago, during the years 1842–1846; together with an excursion into the interior of the eastern part of Java*. Volume 2. T. & W. Boone, London. pp. 335–338.