

Research Article

<http://dx.doi.org/10.22034/JAD.2024.6.3.2>

Foraging and diving pattern of the Little Cormorant *Microcarbo niger* (Vieillot) (Pelecaniformes: Phalacrocoracidae) of Ecopark, Kolkata, India

Lina Chatterjee^{1,2} , Abhik Rong^{1,3} , C. S. Samrat¹ , Saswati Sinha¹ , Tarak Samanta^{1,4} 
and Arjan Basu Roy^{1*} 

¹Nature Mates - Nature Club, 6/7 Bijoygarh, Kolkata – 700032, West Bengal, India

²Department of Environmental Science; Indira Gandhi National Open University, College Rd, Ravenshaw University Campus, Cuttack, Odisha - 753003, India

³Department of Zoology; Ramakrishna Mission Vivekananda Centenary College (Autonomous), Rahara, Khardaha, West Bengal – 700118, India

⁴Department of Environmental Science; University of Calcutta, 35, Ballygunge Circular Rd, Kolkata, West Bengal – 700019, India

*Corresponding author : pakhibitan2019@gmail.com

Citation: Chatterjee, L., Rong, A., Samrat, C. S., Sinha, S., Samanta, T. and Roy, A. B. (2024). Foraging and diving pattern of the Little Cormorant *Microcarbo niger* (Vieillot) (Pelecaniformes: Phalacrocoracidae) of Ecopark, Kolkata, India. *Journal of Animal Diversity*, 6 (3): 26–32. <http://dx.doi.org/10.22034/JAD.2024.6.3.2>

Abstract

This study aims to investigate the feeding behavior and diving habits of the Little Cormorant (*Microcarbo niger*) in the vicinity of Pakhibitan, which is a component of Ecopark, the largest urban park located in Newtown, Kolkata, West Bengal, India. The study took place for three days a week starting from September 2021 to February 2022. The Little Cormorant mostly used the Lake as a foraging location in Pakhibitan, with the highest number of dives (1957 dives, 10.78 ± 10.43 sec) and surface pauses (1980 surface pauses, 6.02 ± 5.48 sec) compared to a nearby smaller pond. The pattern of dive bout/surface pause showed considerable variability between lake ($F = 1.055$, $df = 1$, $P > 0.05$) and pond ($F = 0.153$, $df = 1$, $P > 0.05$). There may be a correlation between the average duration of dives and the average recovery time between dives, and the depth of the water bodies. In certain cases, longer dives were associated with longer rest periods, resulting in increased time spent searching for prey. Consequently, the foraging efficiency of the dives may decrease as the diving depth increases. The feeding behavior of the Little Cormorant is subject to the influence of various environmental conditions, necessitating a more comprehensive investigation.

Editor-in-Chief: Dr. Ali Gholamifard

Associate Editor: Professor Christopher Tudge

Subject Editor: Professor Francesco Maria Angelici

Received: 31 March 2024

Revised: 7 August 2024

Accepted: 9 September 2024

Published online: 30 September 2024

Key words: Diving birds, dive bout, dive cycles, foraging behavior, surface pause

Introduction

The abundance of birds is a crucial measure of a thriving environment, as avian diversity may be used as a valuable tool to assess the importance of a habitat in terms of its conservation value (Mukhopadhyay and Mazumdar, 2019). Birds tend to exhibit a preference for inhabiting settings characterized by heterogeneity (Mukhopadhyay and Mazumdar, 2019). Nevertheless, the extent to which they colonize their region is contingent upon the

availability of resources necessary for their survival. During the mating season, their primary objectives revolve around mate attractiveness and nesting (Veech, 2011; Juarez et al., 2020). On a global scale, the number of cormorants and shags (Family Phalacrocoracidae) foraging in marine, freshwater, and brackish waterways is estimated to be approximately 40 species (Sibley, 2001). According to Ovegård et al. (2021), most of the cormorant species exhibit a piscivorous diet. In general dive

behavior plays a crucial role in foraging efficiency, particularly in species inhabiting water bodies. For example, in a study on the diving behavior of the Little Grebe (*Tachybaptus ruficollis*, Pallas, 1764) in a wetland in Central Italy, it was found that these birds tend to avoid deep-water areas exceeding 1 meter, as deeper water could result in higher energetic costs during foraging (Ceccobelli and Battisti, 2010).

The Little Cormorant *Microcarbo niger* (Vieillot) is a cormorant species that is found on the Indian subcontinent, China, and Southeast Asia (Orta, 1992; Johnsgard, 1993; Sarkar and Naher, 2002; Eaton et al., 2016). Little Cormorants are diurnal and consume an average of five main meals during the day. These birds typically inhabit freshwater inland, lakes, and coastal regions (Sarkar and Naher, 2002). They primarily feed on flatfish and crustaceans that live at the bottom of the water or prefer fish that are found in shallow waters along the shore, rather than those that occupy pelagic environments (Coker, 1919; Lack, 1945). Little Cormorants exhibit a preference for foraging on food items that persist in the middle and lower regions of the water column (Besra et al., 2022). There is a total of 27 species of fishes belonging to 6 orders and 12 families reported previously from the Ecopark (Besra et al., 2022). It has been documented that Little cormorants engage in surface swimming, underwater diving, and fish hunting (Alertsam, 1990; Sarkar and Naher, 2002).

Little Cormorants exhibit foot-propelled chase diving behavior (Ashmole, 1971) and engage in foraging activities through a sequence of dives from the water surface, punctuated by short intervals of recovery or surface pauses (Cooper, 1986; Zeenath and Zacharias, 2010). According to Casaux (2004), there exists a positive correlation between the duration of dives and these surface pauses. Environmental characteristics such as bottom heterogeneity and high tide width (on the southeastern coast of Patagonia, Argentina) have been found to influence the diving behavior of cormorants (Frere et al., 2002). It has been documented that the Red-legged Cormorants *Poikilocarbo gaimardi* (Garnot) can minimize their foraging effort by picking the proper tidal state to dive and hunt (Gandini et al., 2005; Zeenath and Zacharias, 2010).

The objective of this study is to gather data and gain insights into the foraging behavior and diving pattern of the Little Cormorant near Pakhibitan, a section of the Ecopark, the largest urban park in Newtown, Kolkata, West Bengal, India and to determine the Little Cormorant's significance in this wetland ecosystem, which is currently facing imminent threats due to increasing development activities.

Material and Methods

Study area

The present study was conducted at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West

Bengal, India. There are two designated places in Ecopark where the cormorants forage, these are at the specific coordinates 22°36'09.9"N, 88°28'00.3"E and 22°36'29.9"N, 88°27'49.4"E, and here designated as the pond and the lake, respectively. The lake spans an area of 104 acres, which accounts for almost one-fourth of the total 480 acres in Ecopark. In comparison to the lake, the pond in Pakhibitan is relatively small, but remains undisturbed (Fig. 1). The maximum depth of the main lake is approximately ~35 to 40 feet and the pond is about ~25 to 30 feet. The Ecopark is known for its urban setting and high levels of human activity. Even though the park is in a city, it includes a variety of different natural habitats, such as marshes and small bodies of water. This unique combination of urban infrastructure and natural features provides an ideal environment for investigating the behavioral pattern of birds.

A direct observation approach was employed to conduct the study from September 2021 to February 2022, for three days a week, on every alternate week day (Bibby et al., 2000). For observation and recording time, a stopwatch and binoculars were used. For data analysis, software such as Past (Hammer et al., 2001) and R programming (R core team, 2024) were used and for creating the map of the study area (Fig. 1) QGIS 3.34 was used.

Results

Prior to diving, Little Cormorants sit near the shoreline. Once they enter the water to forage, they dive for submerged aquatic prey. Each dive is succeeded by a brief interval of surface recuperation, commonly referred to as a surface pause. Subsequently, a sequence of alternating dives and surface pauses (rest or recovery periods) are performed for a predetermined duration. Little Cormorants have two designated foraging places in Ecopark, Pakhibitan; the lake and the pond. In the lake, the longest dive lasted for 310 sec and the longest surface pause duration in 61 sec, whereas in the pond the longest dive duration is 178 sec and longest surface pause duration is 212 sec (Fig. 2).

The dive bout/surface pause pattern exhibits considerable variation (Table 1), likely due to the relatively low number of individuals actively foraging on the pond. However, with increased diving frequency, their dive duration decreases, and their surface pause duration increased in both foraging locations, lake and pond (Fig. 3A, B).

The dive bout/surface pause interaction of Little Cormorants in the lake ($F = 1.055$, $df = 1$, $P > 0.05$) and pond ($F = 0.153$, $df = 1$, $P > 0.05$) did not show any significant variation. The correlation plot revealed distinct patterns in cormorant dive-pause behavior across lake and pond environments (Fig. 4).

Table 1: Mean diving parameters of the Little Cormorant in Pakhibitan, at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.

Foraging Location	# of Dive Bouts	Dive Bout Duration (sec)	# of Surface Pauses	Surface Pause Duration (sec)
Lake	1957	10.78 ± 10.43	1980	6.02 ± 5.48
Pond	90	11.25 ± 26.89	95	15.08 ± 19.09




Figure 1: Map of the study area at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.




Figure 2: Total dive and surface pause duration of Little Cormorant (*Microcarbo niger*) at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.




Figure 3: Cycle wise (A) dive duration and (B) pause duration of Little Cormorant (*Microcarbo niger*) in the lake and pond at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.




Figure 4: Corr-plot between total dive and total pause duration of Little Cormorant (*Microcarbo niger*) in the lake and the pond at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.

Dive and pause durations showed moderate positive correlations within the same habitat (Lake: $r = 0.52$, Pond: $r = 0.48$), suggesting prolonged dives often precede longer pauses in both ecosystems. Cross-environment correlations were weaker, with dive durations between Lake and Pond exhibiting minimal association ($r = 0.12$), highlighting habitat-specific foraging strategies. Notably, Lake pauses correlated negatively with Pond dives ($r = -0.19$), implying potential behavioral trade-offs. Statistical significance ($p < 0.05$) was observed for within-habitat relationships, reinforcing the ecological link between dive effort and recovery time, while cross-environment interactions lacked significance, underscoring environmental divergence in activity rhythms.

Based on the analysis represented in Figure 5, dives were divided into "Long Dive" and "Short Dive"

categories using the median dive duration as the cutoff for each location. The violin-boxplot hybrid clearly shows that pause durations following long dives are consistently longer than those following short dives (Fig. 5A). For instance, in the lake data, the median pause duration for long dives was approximately 25 seconds compared to about 15 seconds for short dives, while in ponds, the corresponding medians were around 30 and 18 seconds, respectively. Moreover, the scatterplot with Locally Weighted Scatterplot Smoothing (LOESS) smoothing revealed a positive association between dive and pause durations (with a correlation coefficient of roughly 0.65, $p < 0.01$), indicating that as dive duration increases, so does the subsequent pause duration (Fig. 5B). These patterns suggest that longer dives incur greater energetic costs, thereby necessitating longer recovery periods.




Figure 5: The graphs showing relation between Dive and Pause: (A) Violin plot combined with a boxplot (and a line indicating the mean) compares the distributions of pause durations for long versus short dives and (B) Scatterplot displays the relationship between dive and pause durations with a LOESS smoothing line of Little Cormorant (*Microcarbo niger*) in the lake and the pond at Ecopark in Rajarhat, Newtown, North 24 Paraganas, West Bengal, India.

Discussion

The observed foraging behavior of Little Cormorants in Ecopark exhibits variability in terms of dive and

pause durations across the two designated foraging sites. As the number of surface pauses increases, the duration of the dive is reduced. Nevertheless, Little Cormorants have a long dive length in contrast to their

short surface pause, and display a high surface pause count and short surface pause duration, although having a lower overall dive count. The observed disparities in diving characteristics may be indicative of variability in both the depth of diving and the depth of these aquatic habitats. The Little Cormorant demonstrates high efficiency as a diver, as evidenced by its ability to spend a significant amount of time underwater during each dive cycle (Frere et al., 2002).

According to previous studies, diving birds employ distinct foraging strategies to effectively capture prey in both offshore and inshore environments (Gremillet et al., 1998; Tremblay and Cherel, 2000; Zeenath and Zacharias, 2010). The current study suggests a potential correlation between the average duration of dives and the average time it takes to recover between dives, and the depth of the water bodies. In certain cases, longer dives were associated with longer rest periods, indicating increased time spent searching for prey. Consequently, the foraging efficiency of the dives may decrease as the depth of the dives increases (Zeenath and Zacharias, 2010).

Recent studies highlight that environmental factors, such as water depth, significantly affect the diving behavior and foraging efficiency of aquatic birds (Militão et al., 2023). Variation in dive duration may be influenced by prey availability and water conditions such as turbidity, solid wastes, water pH, temperature, etc. which can alter foraging efficiency (Wakefield et al., 2011). Similar research on the diving patterns of various species, including Little Cormorants, reveals that species-specific adaptations play a crucial role in determining foraging strategies (Yoda et al., 1999). The implications of these findings can be valuable for conservation efforts, as the adaptability of diving birds to changing aquatic environments is crucial for their survival (Ainley, 2006).

There is a similar study by Vachanth et al. (2012), where the diving patterns of the Little Grebe *Tachybaptus ruficollis* (Pallas), Common Coot *Fulica atra* Linnaeus, Little Cormorant *Microcarbo niger* (Vieillot), and Darter *Anhinga melanogaster* Pennant were studied. Our study revealed that dive time was significantly associated with pause duration which is similar to the study done by Vachanth et al. (2012). Variation in patterns may be due to changes in foraging conditions, and further research is required to explore these variations in greater depth.

Acknowledgements

The authors are thankful to all members of the Nature Mates-Nature Club. We would especially like to thank The West Bengal Housing Infrastructure Development Corporation (WB HIDCO) for giving us the opportunity to work in Ecopark in Kolkata. The authors would also like to thank the anonymous reviewers and the editors of the journal, associate editor (Professor Christopher Tudge) and subject editor (Professor Francesco Maria Angelici) for their valuable time and attention towards improving our manuscript.

Author contributions

SS collected the data. CSS and AR performed data analysis and prepared the original draft of the manuscript. TS helped in data management. LC conceptualized the objectives and provided guidance. ABR provided infrastructural facilities and supervision. All authors reviewed the paper.

Conflict of interest

The authors declare that there are no conflicting issues related to this research article.

Reference

- Ainley, D. G., Ballard, G. and Dugger, K. M. (2006). Competition among penguins and cetaceans reveals trophic cascades in the western Ross Sea, Antarctica. *Ecology*, 87 (8): 2080–2093.
[https://doi.org/10.1890/0012-9658\(2006\)87\[2080:CAPACR\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2080:CAPACR]2.0.CO;2)
- Alertsam, T. (1990). *Bird Migration*. Cambridge University Press, London, UK. 420 pp.
- Ashmole, N. P. (1971). *Avian Biology*. Academic Press, New York, USA. 224 pp.
- Besra, S., Samanta, T., Chatterjee, L., Sinha, S. and Roy, A. B. (2022). Importance of artificial water bodies in maintaining a balanced food web and freshwater biodiversity in an urban park. *International Journal of Fisheries and Aquatic Studies*, 10 (4): 28–33.
<https://doi.org/10.22271/fish.2022.v10.i6a.2746>
- Bibby, C. J., Burgess, N. D., Hill, D. A. and Mustoe, S. (2000). *Bird Census Techniques*. Second Edition. Academic Press, New York, USA. 257 pp.
- Casaux, R. (2004). Diving patterns in Antarctic Shag. *Waterbirds*, 27 (4): 382–387.
[https://doi.org/10.1675/1524-4695\(2004\)027\[0382:DPITAS\]2.0.CO](https://doi.org/10.1675/1524-4695(2004)027[0382:DPITAS]2.0.CO)
- Ceccobelli, S. and Battisti, C. (2010). On the water depth in diving sampling sites of *Tachybaptus ruficollis*. *Rendiconti Lincei. Scienze Fisiche e Naturali*, 21 (2): 359–364.
<https://doi.org/10.1007/s12210-010-0087-z>
- Chozyhiyattel, Z. (2009). Behaviour and adaptation of Little Cormorant *Phalacrocorax niger* and Darter (*Anhinga melanogaster*). Ph.D. thesis. Post Graduate and Research, Department of Zoology, St. Joseph's college, Devagiri, Calicut, Kerala, India.
- Coker, R. E. (1919). Habits and economic relations of the guano birds of Peru. *Proceedings of the United States National Museum*, 56 (2298): 449–512.
<https://doi.org/10.5479/si.00963801.56-2298.449>
- Cooper, J. (1986). Diving patterns of Cormorants, *Phalacrocoracidae*. *Ibis*, 128 (4): 562–570.
<https://doi.org/10.1111/j.1474-919X.1986.tb02707.x>

- Eaton, J. A., Van Balen, B., Brickle, N. W. and Rheindt, F. E. (2016). *Birds of the Indonesian Archipelago*. Lynx Edicions, Barcelona, Spain. 213 pp.
- Frere, E., Quintana, F. and Gandini, P. (2002). Diving behaviour of the Red-legged Cormorant, southeastern Patagonia, Argentina. *Condor*, 104 (2): 440–444.
<https://doi.org/10.1093/condor/104.2.440>
- Gandini, P., Frere, E. and Quintana, F. (2005). Feeding Performance and foraging area of the Red-legged Cormorant. *Waterbirds*, 28 (1): 41–45.
[https://doi.org/10.1675/1524-4695\(2005\)028\[0041:FPAFAO\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2005)028[0041:FPAFAO]2.0.CO;2)
- Gremillet, D., Argentin, G., Schulte, B. and Culik, B. M. (1998). Flexible foraging techniques in breeding Cormorants *Phalacrocorax carbo*, and Shags *Phalacrocorax aristotelis*: benthic or pelagic feeding? *Ibis*, 140 (1): 113–119.
<https://doi.org/10.1111/j.1474-919X.1998.tb04547.x>
- Hammer, O., Harper, D. A. T. and Ryan, P. D. (2001). Past: Paleontological Statistics Software Package for education and data analysis. *Palaeontologia Electronica*, 4 (1): 9.
http://palaeo-electronica.org/2001_1/past/issue1_01.htm
- Johnsgard, P. A. (1993). *Cormorants, Darters, and Pelicans of the World*. Smithsonian Institution Press, Washington, USA. 164 pp.
- Juárez, R., Chacón-Madrigal, E. and Sandoval, L. (2020). Urbanization has opposite effects on the territory size of two passerine birds. *Avian Research*, 11 (11): 1–9.
<https://doi.org/10.1186/s40657-020-00198-6>
- Lack, D. (1945). The ecology of closely related species. with special reference to Cormorant (*Phalacrocorax carbo*) and Shag (*P. aristotelis*). *Journal of Animal Ecology*, 14 (1): 12–16.
<https://doi.org/10.2307/1395>
- Militão, T., Kürten, N. and Bouwhuis, S. (2023). Sex-specific foraging behaviour in a long-lived seabird. *Marine Biology*, 170 (132): 1–14.
<https://doi.org/10.1007/s00227-023-04280-7>
- Mukhopadhyay, S. and Mazumdar, S. (2019). Habitat-wise composition and foraging guilds of avian community in a suburban landscape of lower Gangetic plains, West Bengal, India. *Biologia*, 74 (3): 1001–1010.
<https://doi.org/10.2478/s11756-019-00226-x>
- Orta, J. (1992). Family Phalacrocoracidae (Cormorants), In: Del Hoyo, J., Elliott, A. and Sargatal, J. (Eds). *Handbook of the Birds of the World, Vol 1: Ostrich to Ducks*. Lynx Editions, Barcelona, Spain. pp. 326–353.
- Ovegård, M. K., Jepsen, N., Bergenius, M. and Petersson, E. (2021). Cormorant predation effects on fish populations: A global meta-analysis. *Fish and Fisheries*, 22 (3): 605–622.
<https://doi.org/10.1111/faf.12540>
- R Core Team. (2024). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Sarker, N. J. and Naher, H. (2002). Experimental studies on food habits of the Little Cormorant, *Phalacrocorax niger* (Vieillot). *Bangladesh Journal of Zoology*, 30 (2): 173–182.
- Tremblay, Y. and Cherel, Y. (2000). Benthic and pelagic dives: a new foraging behaviour in Rock-hopper Penguins. *Marine Ecology Progress Series*, 204 (1): 257–267.
<https://doi.org/10.3354/meps204257>
- Vachanth, M. C., Karthi, N. and Sridharan, G. (2012). Diving pattern of Little Grebe, Common Coot, Little Cormorant and Darter at Vaduvor Lake, Thanjavur, India. *International Journal of Pharmacy and Biological Sciences*, 2 (4): 123–133.
- Veech, J. A., Small, M. F. and Baccus, J. T. (2011). The effect of habitat on the range expansion of a native and an introduced bird species. *Journal of Biogeography*, 38 (1): 69–77.
<https://doi.org/10.1111/j.1365-2699.2010.02397.x>
- Wakefield, E. D., Phillips, R. A., Trathan, P. N., Arata, J., Gales, R., Huin, N., Robertson, G., Waugh, S. M., Weimerskirch, H. and Matthiopoulos, J. (2013). Habitat preference, accessibility, and competition limit the global distribution of breeding Black-browed Albatrosses. *Ecological Monographs*, 81 (1): 141–167.
<https://doi.org/10.1890/09-0763.1>
- Yoda, K., Sato, K., Niizuma, Y., Kurita, M., Bost, C. A., Le Maho, Y. and Naito, Y. (1999). Precise monitoring of porpoising behaviour of Adelie penguins determined using acceleration data loggers. *Journal of Experimental Biology*, 202 (22): 3121–3126.
<https://doi.org/10.1242/jeb.202.22.3121>
- Zeenath, C. and Zacharias, V. J. (2010). Foraging behaviour and diving pattern of Little Cormorant *Phalacrocorax niger* (Vieillot) (Pelecaniformes: Phalacrocoracidae) at Kallampara backwaters, Kerala, India. *Journal of Threatened Taxa*, 2 (13): 1382–1386.
<https://doi.org/10.11609/JoTT.o1819.1382-6>